

ONION				
Sr. no.	Growth Stages	Products	Applications	Method & Dosage
1	Developmental stage	Thrive	For management of fungal disease infection.	Drenching - 2ml/L of water
		Lumme	For healthy growth of plants	Drenching - 2ml/L of water
		Humic acid	For root development	Soil Application - 2gm/ L of water
2	After 45 DAYS OF PLANTATION	Entice	To minimise infestation of insect pest	Drenching or Foliar spray, 2 ml/litre of water
		Spotless	To minimise infestation of bacterial and insect pest	Foliar spray, 2 ml/litre of water
		Lumme	For healthy growth of the bulb	Drenching - 2ml/L of water
		Humic acid	For root development	Soil Application - 2gm/ L of water
		Thrive	For management of fungal disease infection.	Drenching or Foliar spray, 2 ml/litre of water
		Entice	To minimise infestation of insect pest	Drenching or Foliar spray, 2 ml/litre of water
<p>Note:</p> <ol style="list-style-type: none"> 1. Spray should be given in morning or evening. Avoid applications under high wind, there should not be any heavy rainfall at least within 5 hours of application. 2. Not to be used on crops other than specified on this leaflet. 3. Do not mix above Herbal products with any chemical products. 4. The used packages shall not be left outside to prevent their re-use. This material should be disposed of in safe manner to prevent environmental and water pollution. 				